

Classification of degree three arcs in $PG(2,19)$ up to stabilizer groups

Islam A. Abd-alzahra, Mohammed A. Ibrahim *

Department of Mathematics, College of Science, University of Basrah, Basra, Iraq.

*Corresponding author E-mail: mohammed.ibrahim@uobasrah.edu.iq

<https://doi.org/10.29072/basjs.20240201>

ARTICLE INFO	ABSTRACT
<p>Keywords Projective Plane; Arcs ; Group action; Stabilizer Group</p>	<p>An $(n, 3)$-arc \mathcal{K} in projective plane $PG(2, q)$ of size n and degree three is a set of n points such that every line in the plane meet it in less than or equal three points, also the arc \mathcal{K} is complete if it is not contained in $(n + 1, 3)$-arc. In this paper, the classification of degree three arcs in $PG(2, 19)$ is introduced in details according to their stabilizer groups. The motivation for working in the projective plane of order 19 is twofold. First, the size of the largest $(n, 3)$-arc is not known. Second, the number of $(n, 3)$-arcs is significantly higher in the projective plane of order 19 than it is in the projective plane of order q for $q < 19$, giving a large number of $(n, 3)$-arcs for the study.</p>

Received 27 Apr 2024; Received in revised form 2 July 2024; Accepted 4 Aug 2024, Published 30 Aug 2024



1. Introduction

Let \mathbb{F}_q be the Galois field with q elements, and $V(3, q)$ be the 3-dimensional vector space over the field \mathbb{F}_q . The corresponding projective plane of $V(3, q)$ is denoted by \mathbb{P}_q^2 . The points (x_1, x_2, x_3) of \mathbb{P}_q^2 are the 1-dimensional subspaces of $V(3, q)$. Subspaces of dimension two of $V(3, q)$ are called lines which has the form $V(ax_1 + bx_2 + cx_3)$. The number of points and the number of lines in \mathbb{P}_q^2 is $q^2 + q + 1$. There are $q + 1$ points on every line and $q + 1$ lines through every point. Many researches have been studies the subject of projective geometries for examples see [1], the tools of this paper Gap-Groups, Algorithms, programming a system for computational discrete algebra [17].

A full classification of $(n, 3)$ -arcs are given by Cook [2]. For $q = 11$, a maximal $(n, 3)$ -arcs has been found by Marcugini [3]. For $q = 13$, the maximal arc has been found in [4]. Many studies and improvements have been given to get a largest (n, r) -arc of size two, three, four, etc. For more detailed to the size of $(n, 3)$ -arcs see [5]. A full classification of the complete k -arcs of $PG(2, 23)$ and $PG(2, 25)$ is done by Coolsaet and Sticker [6]. Also, the complete $(k, 3)$ -arcs of $PG(2, q)$, $q \leq 13$ is given by Coolsaet and Sticker [7]. The classification of the $(k, 3)$ -arcs in $PG(2, 37)$ is presented [8]. In [9], the construction of $(4v, 3)$ -arcs that produced from irreducible plane-cubic is discussed for all values of q , where $7 \leq q \leq 37$. Many large complete arcs such as $(46, 3)$ -arcs, $(67, 4)$ -arcs, $(91, 5)$ -arcs, $(201, 8)$ -arcs, $(226, 9)$ -arcs, $(469, 16)$ -arcs and $(488, 17)$ -arcs in $PG(2, 37)$ are obtained in [10]. Furthermore, large size for the complete $(k, 3)$ -arcs in the projective plane of order nineteen $PG(2, 19)$ using the method of secants distributions [11].

The main aim of this paper is to determine the stabilizer group of each an equivalent $(n, 3)$ -arcs in \mathbb{P}_q^2 , where $n > 4$. Classify $(n, 3)$ -arcs in way, give us $((n, 3)$ -arcs of large size in \mathbb{P}_{19}^2 . The classification technique is based on constructed the special arcs in \mathbb{P}_q^2 to find the largest size of complete $(n, 3)$ -arcs [12]. In fact, up to stabilizer group we obtained that the largest size of complete $(n, 3)$ -arc is equal to 30.

2. Preliminaries

2.2 Group Theory

A group is a set G together with an operation $*$ satisfying the following requirements:

1. for each pair (x, y) of elements of $G \times G$, $x * y$ is an element of G ;



2. for all elements x, y, z of G , $(x * y) * z = x * (y * z)$;
3. there is an element $e \in G$ such that $e * g = g = g * e$ for all $g \in G$;
4. given an element $g \in G$, there is an element $g' \in G$ such that $g' * g = e = g * g'$.

A group G is *abelian* if, for all $g, g' \in G$, $g' * g = g * g'$. A *cyclic group* is a group generated by a single element; that is, a group consisting of all powers of one of its elements. If G is a cyclic group generated by $g \in G$, we write $G = \langle g \rangle$. A finite group G of order n is cyclic if and only if it contains an element of order n . Also, a group of prime order is cyclic [13].

Definition ([13]) An *action* of a group $(G, *)$ on a set K is a function $\varphi : K \times G \rightarrow K$ with the following properties:

1. $((x, g)\varphi, h)\varphi = (x, g * h)\varphi$ for all $x \in K$ and $g, h \in G$;
2. $(x, e)\varphi = x$ for all $x \in K$, where e is the identity of G ;
3. $((x, g)\varphi, g')\mu = ((x, g')\varphi, g)\varphi = x$ for all $x \in K$; $g \in G$, where g' is the inverse of the element g in G .

Definition ([13]) Let G, H be two groups. A *homomorphism* $h : G \rightarrow H$ is a function h from G to H that satisfies the condition, for all $g_1, g_2 \in G$,

$$(g_1 g_2)h = (g_1 h)(g_2 h) \quad (1)$$

A homomorphism that is one-to-one and onto is called an *isomorphism*. In this case, G and H is called *isomorphic* groups. A bijective homomorphism h from a group to itself is called an *automorphism*.

Definition ([14]) A bijection $h : X \rightarrow X$ is a permutation on X . The set of all permutations on X is denoted by $S(X)$.

The set $S(X)$ forms a group under the usual composition of functions. If $X = \{1, 2, \dots, n\}$ or any set with cardinality n , then $S(X)$ is written as \mathfrak{S}_n which is called the symmetric group on n symbols.

2.2 Projective Spaces Over a Finite Field

A field is a set \mathbb{F} closed under two operations $+, \times$ such that

1. $(\mathbb{F}, +)$ is an abelian group with identity 0 ;



2. $(\mathbb{F} \setminus \{0\}, \times)$ is an abelian group with identity 1;
3. $a(b + c) = ab + ac$; $(a + b)c = ac + bc$, for all $a, b, c \in \mathbb{F}$.

Let \mathbb{F}_q denotes a field of q elements. Let $V = V(n + 1, q)$ be an $(n + 1)$ -dimensional vector space over a field \mathbb{F}_q with zero element $\mathbf{0}$. Consider the equivalence relation on the elements of $V^* = V \setminus \{\mathbf{0}\}$ whose equivalence classes are the 1-dimensional subspaces of V with zero removed. In fact, if $v_1, v_2 \in V^*$ where $v_1 = (x_1, \dots, x_{n+1})$ and $v_2 = (y_1, \dots, y_{n+1})$, then v_1 is equivalent to v_2 if $v_2 = \alpha v_1$ for some $\alpha \in \mathbb{F}_q \setminus \{0\}$; that is, $y_i = \alpha x_i$ for all i . The set of all equivalence classes is the n -dimensional projective space over \mathbb{F}_q and is denoted by $PG(n, q)$.

The elements of $PG(n, q)$ are called *points*; the equivalence class of the vector v is the point $P(v)$. In this case, we say that v is a *coordinate vector* for $P(v)$ or that v is a vector representing $P(v)$. By definition, $P(\alpha v) = P(v)$ for all $\alpha \in \mathbb{F}_q \setminus \{0\}$.

Definition ([15]) A collineation of projective plane $\Pi = PG(2, q)$ is a map φ of Π onto Π such that φ is bijection;

1. φ maps points onto points and lines onto lines;
2. if P and ℓ are an incident point and line in Π , then $\varphi(P)$ and $\varphi(\ell)$ are incident.

Theorem ([15]) If $\{P_0, P_1, \dots, P_{n+1}\}, \{P'_0, P'_1, \dots, P'_{n+1}\}$ are two sets of $n + 2$ points of $PG(n, q)$ such that no $n + 1$ points chosen from the same set lie in a subspace of dimension $n - 1$, then there exists a unique projectivity \mathfrak{Z} such that $P'_i = P_i \mathfrak{Z}$, for all $i = 0, 1, \dots, n + 1$.

The Theorem above is called the Fundamental Theorem of Projective Geometry.

Definition ([15]) For any positive integer n , the *general linear group*, $GL(n, q)$, is the set of all invertible $n \times n$ matrices over \mathbb{F}_q under matrix multiplication. The order of the group $GL(n, q)$ is $(q^n - 1)(q^n - q)(q^n - q^2) \cdots (q^n - q^{n-1})$.

The *projective general linear group* $PGL(n, q)$ is the group of projectivities of $PG(n - 1, q)$ with respect to the operation of composition of maps.

Definition ([15]) The *special linear group* $SL(n, q)$ is the subgroup of the group $GL(n, q)$ consisting of all non-singular matrices. The *projective special linear group* $PSL(n, q)$ is the quotient group $SL(n, q)/Z$, where Z is the subgroup of scalar matrices in $SL(n, q)$.



2.3 Projective Planes

Consider the projective plane $PG(2, q)$ over the field \mathbb{F}_q . The projective plane $PG(2, q)$ contains $q^2 + q + 1$ points and $q^2 + q + 1$ lines. There are exactly $q + 1$ points on each line, and $q + 1$ lines through each point. The points and lines of $PG(2, q)$ satisfy the following axioms of a projective plane:

1. every two distinct points are on a unique common line;
2. every two distinct lines contain a unique common point;
3. there are four distinct points, no three of which are on a common line.

A point $P(x, y, z)$ is incident with a line $\ell(k, l, m)$ if and only if $kx + ly + mz = 0$.

There is a non-singular matrix $\mathfrak{T} = (\alpha_{ij})$ where $\alpha_{ij} \in \mathbb{F}_q$ associated to each bijection from $PG(2, q)$ to $PG(2, q)$ that maps the point $P(v) = P(x, y, z)$ to the point $P(v') = P(x', y', z')$ and the line $\ell(U) = \ell(k, l, m)$ to the line $\ell(U'^t) = \ell(k', l', m')^t$ with $v' = v\mathfrak{T}$ and $U'^t = \mathfrak{T}^{-1}U^t$.

In other words,

$$(x', y', z') = (x, y, z) \begin{bmatrix} \alpha_{11} & \alpha_{12} & \alpha_{13} \\ \alpha_{21} & \alpha_{22} & \alpha_{23} \\ \alpha_{31} & \alpha_{32} & \alpha_{33} \end{bmatrix} \quad (2)$$

and

$$\begin{bmatrix} k' \\ l' \\ m' \end{bmatrix} = \begin{bmatrix} \alpha_{11} & \alpha_{12} & \alpha_{13} \\ \alpha_{21} & \alpha_{22} & \alpha_{23} \\ \alpha_{31} & \alpha_{32} & \alpha_{33} \end{bmatrix}^{-1} \begin{bmatrix} k \\ l \\ m \end{bmatrix} \quad (3)$$

Any such a bijection is called a *projectivity* or *projective linear transformation*. Further, a projectivity preserves the incidence between points and lines.

The group of all projectivities of $PG(2, q)$ is the projective general linear group $PGL(3, q)$ and has order $(q^3 - 1)(q^2 - 1)q^3$. From the Fundamental Theorem of Projective Geometry, a projectivity is uniquely determined by the four images of the vertices of a quadrangle.

Definition ([13]) A (k, n) -arc \mathcal{K} in a projective plane $PG(2, q)$ is a set of k points such that some line of the plane meets \mathcal{K} in n points but such that no line meets \mathcal{K} in more than n points, where $n \geq 2$.



Throughout, \mathbb{P}_q^2 will denote the projective plane $PG(2, q)$. A line ℓ of \mathbb{P}_q^2 is an i -secant of a (k, n) -arc \mathcal{K} if ℓ intersects \mathcal{K} in i points. Let τ_i be the total number of i -secants to \mathcal{K} . The number of i -secants to \mathcal{K} through a point P of \mathcal{K} is denoted by ρ_i or $\rho_i(P)$. Moreover, σ_i or $\sigma_i(Q)$ denotes the number of i -secants to \mathcal{K} through a point Q of $\mathbb{P}_q^2 \setminus \mathcal{K}$. A (k, n) -arc is complete if there is no $(k + 1, n)$ -arc containing it. For more information about complete and incomplete (k, n) -arcs, one can see [16].

Lemma ([15]) For a (k, n) -arc \mathcal{K} , the following equations hold:

$$\sum_{i=0}^n \tau_i = q^2 + q + 1; \tag{4}$$

$$\sum_{i=1}^n i\tau_i = k(q + 1); \tag{5}$$

$$\sum_{i=2}^n i(i - 1)\tau_i = k(k - 1); \tag{6}$$

Definition ([10]) The points out of a (k, n) -arc \mathcal{K} in \mathbb{P}_q^2 which passes through it i -secant of \mathcal{K} is called a point of index i .

1. Main Results

3.1 The classification of $(\nu, 3)$ -arcs; $\nu = 5, 6, 7$

First of all, let us give the following definitions which help us in our study.

Definition ([9]). Two (k, n) -arcs \mathcal{K}_1 and \mathcal{K}_2 in \mathbb{P}_q^2 are said to be stabilizer inequivalent if they have different stabilizer groups, that is, $\text{Stab}(\mathcal{K}_1) \not\cong \text{Stab}(\mathcal{K}_2)$ where

$$\text{Stab}(\mathcal{K}) = \{\mathfrak{T} \in PGL(3, q) : \mathfrak{T}(\mathcal{K}) = \mathcal{K}\},$$

for any (k, n) -arc \mathcal{K} in \mathbb{P}_q^2 .

Definition. A point P in \mathbb{P}_q^2 is called preferred point of type 3 if when it added to ν -arc gives an arc of degree 3 in \mathbb{P}_q^2 .



Let $\mathcal{R} = \{1,2,3,4\}$ be the 2-arc consisting of the indices of the points $P_1 = (1,0,0)$, $P_2 = (0,1,0)$, $P_3 = (0,0,1)$ and $P_4 = (1,1,1)$. Henceforth, \mathcal{D} denotes a set of preferred points of type 3 in \mathbb{P}_q^2 which can be add to $\mathcal{R} = \{1,2,3,4\}$ to produce arc of degree three.

In this subsection, the classification of $(v, 3)$ -arcs, where $v = 5,6,7$, is established by classifying all such arcs up to stabilizer groups.

Let $\mathcal{R} = \{1,2,3,4\}$ be the 2-arc consisting of the indices of the points $P_1 = (1,0,0)$, $P_2 = (0,1,0)$, $P_3 = (0,0,1)$ and $P_4 = (1,1,1)$. There is only one inequivalent $(4,3)$ -arc in \mathbb{P}_{19}^2 up to stabilizer group, while there are 4 types of $(5,3)$ -arcs during implementation our program. The 4 types are shown in Table 1.

Table 1. Types of $(5, 3)$ -arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_5^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_5^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_5^j)$
\mathcal{A}_5^1	5	\mathbb{Z}_2
\mathcal{A}_5^2	35	$\mathbb{Z}_2 \times \mathbb{Z}_2$
\mathcal{A}_5^3	25	\mathbb{Z}_6
\mathcal{A}_5^4	150	\mathbf{D}_4

Note that, $\text{Stab}(\mathcal{A}_5^1) = \langle M_5^1 \rangle$ and $\text{Stab}(\mathcal{A}_5^3) = \langle M_5^3 \rangle$, where

$$M_5^1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \text{ and } M_5^3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$$

The stabilizer group \mathcal{A}_5^2 consists of the following 4 projective matrices each one of them has order 2, say

$$M_{5,1}^2 = I, M_{5,2}^2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}, M_{5,3}^2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ and } M_{5,4}^2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$$



Moreover, the group $\text{Stab}(\mathcal{A}_5^3)$ is cyclic of order 6 and $\text{Stab}(\mathcal{A}_5^3) = \langle M_5^3 \rangle$, where

$$M_5^3 = \begin{bmatrix} 0 & -1 & 1 \\ 7 & -8 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

The stabilizer group of \mathcal{A}_5^4 consists of the following 8 projective matrices, say

$$M_{5,1}^4 = I, M_{5,2}^4 = \begin{bmatrix} 0 & 0 & 1 \\ -1 & 0 & 1 \\ 0 & -1 & 1 \end{bmatrix}, M_{5,3}^4 = \begin{bmatrix} 0 & 1 & -1 \\ 0 & 1 & 0 \\ -1 & 1 & 0 \end{bmatrix}, M_{5,4}^4 = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 0 & -1 \\ 1 & 0 & 0 \end{bmatrix},$$

$$M_{5,5}^4 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}, M_{5,6}^4 = \begin{bmatrix} 0 & -1 & 1 \\ -1 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}, M_{5,7}^4 = \begin{bmatrix} -1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & -1 \end{bmatrix} \text{ and } M_{5,8}^4 = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & -1 \\ 1 & -1 & 0 \end{bmatrix}.$$

The order of $M_{5,1}^4$ is 1, the order of both $M_{5,2}^4, M_{5,4}^4$ is 4, and the order of each matrix in the remaining projective matrices is 2. So, $\text{Stab}(\mathcal{A}_5^3)$ isomorphic to \mathbf{D}_4 .

Now, by adding 2 preferred points of type 3 to \mathcal{R} , we get (6,3)-arcs. In fact, there are 8 types of (6,3)-arcs up to stabilizer group, as shown in Table 2.

Table 2. Types of (6, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_6^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_6^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_6^j)$
\mathcal{A}_6^1	5,6	\mathbf{I}
\mathcal{A}_6^2	5,12	\mathbb{Z}_2
\mathcal{A}_6^3	5,164	\mathbb{Z}_3
\mathcal{A}_6^4	35,56	\mathfrak{S}_3
\mathcal{A}_6^5	35,89	\mathbf{D}_4



\mathcal{A}_6^6	25,133	$\mathfrak{S}_3 \times \mathbb{Z}_3$
\mathcal{A}_6^7	150,236	\mathfrak{S}_4
\mathcal{A}_6^8	40,89	\mathbb{Z}_6

In Table 2, $\text{Stab}(\mathcal{A}_6^2) = \langle M_6^2 \rangle$, $\text{Stab}(\mathcal{A}_6^3) = \langle M_6^3 \rangle$ and $\text{Stab}(\mathcal{A}_6^8) = \langle M_6^8 \rangle$, where

$$M_6^2 = \begin{bmatrix} -4 & 4 & 0 \\ -3 & 4 & 0 \\ -6 & 4 & 2 \end{bmatrix}, M_6^3 = \begin{bmatrix} -5 & 6 & 0 \\ -5 & 5 & -8 \\ -5 & -3 & 0 \end{bmatrix} \text{ and } M_6^8 = \begin{bmatrix} 7 & 8 & 5 \\ 0 & 8 & 0 \\ 0 & 8 & -7 \end{bmatrix}.$$

The group $\text{Stab}(\mathcal{A}_6^4)$ is non-abelian of order 6. The following are all the projective matrices in $\text{Stab}(\mathcal{A}_6^4)$:

$$M_{6,1}^4 = I, M_{6,2}^4 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{bmatrix}, M_{6,3}^4 = \begin{bmatrix} -5 & -5 & -9 \\ -4 & 5 & 0 \\ 5 & -5 & 0 \end{bmatrix}, M_{6,4}^4 = \begin{bmatrix} 1 & -9 & 9 \\ 0 & 9 & -8 \\ 0 & -9 & -9 \end{bmatrix},$$

$$M_{6,5}^4 = \begin{bmatrix} -5 & 5 & 0 \\ -4 & 5 & 0 \\ 5 & 5 & 9 \end{bmatrix} \text{ and } M_{6,6}^4 = \begin{bmatrix} 0 & -9 & -9 \\ 0 & -9 & 8 \\ 1 & -9 & 9 \end{bmatrix}.$$

Moreover, $\text{Stab}(\mathcal{A}_6^5) = \langle R, S: R^4 = S^2 = I, SRS^{-1} = R^{-1} \rangle$, where

$$R = \begin{bmatrix} 0 & -1 & 1 \\ 1 & 0 & -1 \\ 0 & 0 & 1 \end{bmatrix} \text{ and } S = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

The group $\text{Stab}(\mathcal{A}_6^6)$ has 18 projective matrices, say

$$M_{6,1}^6 = I, M_{6,2}^6 = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & -1 \\ 1 & -1 & 0 \end{bmatrix}, M_{6,3}^6 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 1 & -1 \\ 7 & -7 & 0 \end{bmatrix}, M_{6,4}^6 = \begin{bmatrix} -8 & 0 & 8 \\ 0 & -8 & 8 \\ 0 & 0 & 1 \end{bmatrix},$$



$$M_{6,5}^6 = \begin{bmatrix} 8 & 0 & -7 \\ 0 & 8 & -7 \\ 0 & 0 & -7 \end{bmatrix}, M_{6,6}^6 = \begin{bmatrix} -7 & 8 & 0 \\ -7 & 8 & -1 \\ -7 & 8 & 0 \end{bmatrix}, M_{6,7}^6 = \begin{bmatrix} -7 & 8 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, M_{6,8}^6 = \begin{bmatrix} 8 & 0 & -7 \\ 8 & 0 & 0 \\ 8 & -8 & 0 \end{bmatrix},$$

$$M_{6,9}^6 = \begin{bmatrix} -1 & -7 & 8 \\ -8 & 0 & 8 \\ 0 & 0 & 1 \end{bmatrix}, M_{6,10}^6 = \begin{bmatrix} 0 & 8 & -7 \\ 0 & 8 & 0 \\ -1 & 1 & 0 \end{bmatrix}, M_{6,11}^6 = \begin{bmatrix} 1 & 7 & -7 \\ 1 & 7 & 0 \\ -7 & 7 & 0 \end{bmatrix},$$

$$M_{6,12}^6 = \begin{bmatrix} 1 & 7 & -7 \\ 8 & 0 & -7 \\ 0 & 0 & -7 \end{bmatrix}, M_{6,13}^6 = \begin{bmatrix} 0 & 1 & 0 \\ -7 & 8 & 0 \\ 0 & 0 & 1 \end{bmatrix}, M_{6,14}^6 = \begin{bmatrix} 8 & 0 & -8 \\ 8 & 0 & -7 \\ 8 & -8 & 0 \end{bmatrix},$$

$$M_{6,15}^6 = \begin{bmatrix} 0 & -8 & 8 \\ -1 & -7 & 8 \\ 0 & 0 & 1 \end{bmatrix}, M_{6,16}^6 = \begin{bmatrix} 0 & 8 & -8 \\ 0 & 8 & -7 \\ -1 & 1 & 0 \end{bmatrix}, M_{6,17}^6 = \begin{bmatrix} 1 & 7 & -8 \\ 1 & 7 & -7 \\ -7 & 7 & 0 \end{bmatrix} \text{ and}$$

$$M_{6,18}^6 = \begin{bmatrix} 0 & 8 & -7 \\ 1 & 7 & -7 \\ 0 & 0 & -7 \end{bmatrix}.$$

The group $\text{Stab}(\mathcal{A}_6^7)$ is non-abelian, and has 24 projective matrices. In fact, $\text{Stab}(\mathcal{A}_6^7) = \langle R, S \rangle$, where

$$R = \begin{bmatrix} 0 & -1 & 1 \\ 0 & 0 & 1 \\ -1 & 0 & 1 \end{bmatrix} \text{ and } S = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$$

Adding 3 preferred points of type 3 to \mathcal{R} , give us (7,3)-arcs. Further, there are 9 types of (7,3)-arcs up to stabilizer group, as shown in Table 3.

Table 3. Types of (7, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_7^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_7^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_7^j)$

\mathcal{A}_7^1	5,6,17	I
\mathcal{A}_7^2	5,6,236	\mathbb{Z}_2
\mathcal{A}_7^3	5,6,336	\mathbb{Z}_3
\mathcal{A}_7^4	9,115,188	\mathbb{Z}_4
\mathcal{A}_7^5	9,98,256	$\mathbb{Z}_2 \times \mathbb{Z}_2$
\mathcal{A}_7^6	9,59,256	\mathfrak{S}_3
\mathcal{A}_7^7	26,285,339	\mathbb{Z}_6
\mathcal{A}_7^8	5,12,344	D₆
\mathcal{A}_7^9	99,128,305	\mathfrak{S}_4

In Table 3, $\text{Stab}(\mathcal{A}_7^2) = \langle M_7^2 \rangle$, $\text{Stab}(\mathcal{A}_7^3) = \langle M_7^3 \rangle$, $\text{Stab}(\mathcal{A}_7^4) = \langle M_7^4 \rangle$ and $\text{Stab}(\mathcal{A}_7^5) = \langle M_7^5 \rangle$, where

$$M_7^2 = \begin{bmatrix} -3 & 9 & -6 \\ 0 & 9 & -8 \\ 0 & 9 & -9 \end{bmatrix}, M_7^3 = \begin{bmatrix} 0 & 1 & 0 \\ -7 & -8 & 0 \\ 0 & -8 & 8 \end{bmatrix}, M_7^4 = \begin{bmatrix} 0 & 6 & -5 \\ 2 & -8 & 0 \\ 0 & -2 & 0 \end{bmatrix} \text{ and } M_7^5 = \begin{bmatrix} 0 & 8 & -8 \\ -1 & 1 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

The stabilizer group of the (7,3)-arc, \mathcal{A}_7^5 consists of the following 4 projective matrices:

$$M_{7,1}^5 = I, M_{7,2}^5 = \begin{bmatrix} 0 & 6 & -5 \\ -3 & 0 & 4 \\ 0 & 0 & 1 \end{bmatrix}, M_{7,3}^5 = \begin{bmatrix} 0 & 6 & -5 \\ 0 & -1 & 0 \\ -4 & -5 & 0 \end{bmatrix} \text{ and } M_{7,4}^5 = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 0 & -4 \\ -4 & -5 & 0 \end{bmatrix}.$$

All the previous matrices has order 2 except the identity matrix. Hence, the stabilizer group of \mathcal{A}_7^5 is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$.

The group $\text{Stab}(\mathcal{A}_7^6)$ is non-abelian generating by the projective matrices R and S , where



$$R = \begin{bmatrix} 0 & 6 & -5 \\ 3 & 6 & -9 \\ 0 & 6 & -6 \end{bmatrix} \text{ and } S = \begin{bmatrix} 1 & 0 & 0 \\ 1 & -1 & 0 \\ 1 & 0 & -1 \end{bmatrix}.$$

In fact, $\text{Stab}(\mathcal{A}_7^6)$ is isomorphic to \mathfrak{S}_3 .

The stabilizer group of the (7,3)-arc, \mathcal{A}_7^8 is non-abelian consisting of the 12 projective matrices, and $\text{Stab}(\mathcal{A}_7^8) = \langle R, S: R^6 = S^2 = I, SRS^{-1} = R^{-1} \rangle$, where

$$R = \begin{bmatrix} 0 & 2 & -1 \\ -7 & 3 & 0 \\ 0 & -8 & 0 \end{bmatrix} \text{ and } S = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$$

Finally, the group of the (7,3)-arc, \mathcal{A}_7^9 has 24 projective matrices, and it is isomorphic to \mathfrak{S}_4 . Moreover, $\text{Stab}(\mathcal{A}_7^9) = \langle R, S \rangle$, where

$$R = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix} \text{ and } S = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$$

3.2 The classification of $(\nu, 3)$ -arcs; $\nu = 8, 9, 10$

In this subsection, the classification of $(\nu, 3)$ -arcs, where $\nu = 8, 9, 10$, is established by classifying all such arcs up to stabilizer groups. By adding 4 preferred points of type 3 to \mathcal{R} , we get the following (8,3)-arcs as illustrated in Table 4.

Table 4. Types of (8, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_8^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_8^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_8^j)$
\mathcal{A}_8^1	5, 6, 7, 8	I
\mathcal{A}_8^2	5, 6, 7, 330	\mathbb{Z}_2
\mathcal{A}_8^3	5, 6, 336, 265	\mathbb{Z}_3



\mathcal{A}_8^4	5, 12, 344, 157	$\mathbb{Z}_2 \times \mathbb{Z}_2$
\mathcal{A}_8^5	5, 12, 344, 282	\mathbb{Z}_6
\mathcal{A}_8^6	25, 133, 40, 237	SL(2, 3)

Let us explain the groups appeared in Table 4.

$\text{Stab}(\mathcal{A}_8^2) = \langle M_8^2 \rangle$, $\text{Stab}(\mathcal{A}_8^3) = \langle M_8^3 \rangle$, and $\text{Stab}(\mathcal{A}_8^5) = \langle M_8^5 \rangle$, where

$$M_8^2 = \begin{bmatrix} 3 & 1 & -3 \\ 0 & -5 & 5 \\ 0 & -7 & 5 \end{bmatrix}, M_8^3 = \begin{bmatrix} 0 & 1 & 0 \\ -7 & -8 & 0 \\ 0 & -8 & 8 \end{bmatrix} \text{ and } M_8^5 = \begin{bmatrix} 0 & 2 & -1 \\ -7 & 3 & 0 \\ 0 & -8 & 0 \end{bmatrix}.$$

The stabilizer group of the (8,3)-arc, \mathcal{A}_8^4 consists of the following 4 projective matrices:

$$M_{8,1}^4 = I, M_{8,2}^4 = \begin{bmatrix} 0 & -1 & 2 \\ -7 & 0 & 3 \\ 0 & 0 & -8 \end{bmatrix}, M_{8,3}^4 = \begin{bmatrix} 9 & -4 & -4 \\ 9 & -3 & -6 \\ 9 & -6 & -3 \end{bmatrix} \text{ and } M_{8,4}^4 = \begin{bmatrix} -4 & 4 & 0 \\ -3 & 4 & 0 \\ -6 & 4 & 2 \end{bmatrix}.$$

All the previous matrices has order 2 except the identity matrix.

The stabilizer group of the (8,3)-arc, \mathcal{A}_8^6 is non-abelian consisting 24 projective matrices. In

fact,

$\text{Stab}(\mathcal{A}_8^6) = \langle A, B, C : A^4 = C^3 = I, A^2 = B^2, BAB^{-1} = A^{-1}, CAC^{-1} = B, CBC^{-1} = AB \rangle$, where

$$A = \begin{bmatrix} 8 & -8 & 0 \\ 8 & 0 & -7 \\ 1 & 0 & 0 \end{bmatrix}, B = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ -8 & 1 & 0 \end{bmatrix} \text{ and } C = \begin{bmatrix} 7 & -7 & 0 \\ 0 & 1 & 0 \\ 0 & -7 & 7 \end{bmatrix}.$$

Now, adding 5 preferred points of type 3 to \mathcal{R} , gives us (9,3)-arcs as shown in Table 5.

Table 5. Types of (9, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_9^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_9^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_9^j)$
-------------------	-------------------------	--------------------------------

\mathcal{A}_9^1	5,6,50,52,376	I
\mathcal{A}_9^2	5,41,43,247,363	\mathbb{Z}_2
\mathcal{A}_9^3	5,6,265,336,357	\mathbb{Z}_3
\mathcal{A}_9^4	5,12,157,303,344	$\mathbb{Z}_2 \times \mathbb{Z}_2$
\mathcal{A}_9^5	25,37,40,133,237	\mathbb{Z}_6
\mathcal{A}_9^6	5,12,282,327,344	D₆
\mathcal{A}_9^7	25,40,133,237,247	G₂₁₆

In Table 5, $\text{Stab}(\mathcal{A}_9^2) = \langle M_9^2 \rangle$, $\text{Stab}(\mathcal{A}_9^3) = \langle M_9^3 \rangle$, and $\text{Stab}(\mathcal{A}_9^5) = \langle M_9^5 \rangle$, where

$$M_9^2 = \begin{bmatrix} 9 & -8 & 0 \\ 9 & -9 & 0 \\ 7 & 8 & -3 \end{bmatrix}, M_9^3 = \begin{bmatrix} 0 & 1 & 0 \\ -7 & -8 & 0 \\ 0 & -8 & 8 \end{bmatrix} \text{ and } M_9^5 = \begin{bmatrix} 1 & -1 & 0 \\ 8 & -1 & -7 \\ 1 & 7 & -7 \end{bmatrix}.$$

The stabilizer group of the (9,3)-arc, \mathcal{A}_9^4 consists of the 4 projective matrices, each one of them has of order 2 except the identity matrix:

$$M_{9,1}^4 = I, M_{9,2}^4 = \begin{bmatrix} 0 & 2 & -1 \\ 0 & -8 & 0 \\ -7 & 3 & 0 \end{bmatrix}, M_{9,3}^4 = \begin{bmatrix} 9 & -4 & -4 \\ 9 & -3 & -6 \\ 9 & -6 & -3 \end{bmatrix} \text{ and } M_{9,4}^4 = \begin{bmatrix} -4 & 0 & 4 \\ -6 & 2 & 4 \\ -3 & 0 & 4 \end{bmatrix}.$$

The stabilizer group of the (9,3)-arc, \mathcal{A}_9^6 is non-abelian consisting of the 12 projective matrices, and $\text{Stab}(\mathcal{A}_9^6) = \langle R, S: R^6 = S^2 = I, SRS^{-1} = R^{-1} \rangle$, where

$$R = \begin{bmatrix} 0 & 2 & -1 \\ -7 & 3 & 0 \\ 0 & -8 & 0 \end{bmatrix} \text{ and } S = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$$

The group, $\text{Stab}(\mathcal{A}_9^7)$, is non-abelian with 216 projective matrices. In fact,

- (1) the identity matrix has order 1, say $M_1 = I$,
- (2) there are 9 projective matrices of order 2, some of them are:



$$M_2 = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & -1 \\ 1 & -1 & 0 \end{bmatrix}, M_3 = \begin{bmatrix} 0 & 8 & -7 \\ -1 & 0 & 1 \\ 0 & 0 & -7 \end{bmatrix}, \dots, M_{10},$$

(3) there are 80 projective matrices of order 3, some of them are:

$$M_{11} = \begin{bmatrix} 7 & -7 & 0 \\ 0 & 1 & 0 \\ 0 & -7 & 7 \end{bmatrix}, M_{12} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 8 \\ 1 & 0 & 0 \end{bmatrix}, \dots, M_{90},$$

(4) there are 54 projective matrices of order 4, some of them are:

$$M_{91} = \begin{bmatrix} 0 & -1 & 1 \\ 0 & 0 & 1 \\ -1 & 0 & 1 \end{bmatrix}, M_{92} = \begin{bmatrix} 0 & 1 & -1 \\ -1 & 1 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \dots, M_{144},$$

(5) there are 72 projective matrices of order 6, some of them are:

$$M_{145} = \begin{bmatrix} 0 & 0 & 1 \\ -8 & 0 & 8 \\ 0 & -1 & 1 \end{bmatrix}, M_{146} = \begin{bmatrix} 1 & 0 & -1 \\ 8 & -8 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \dots, M_{216}.$$

Adding 6 preferred points of type 3 to \mathcal{R} , gives us six (10,3)-arcs up to stabilizer group as illustrated in Table 6.

Table 6. Types of (10, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{10}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{10}^j	Points of \mathcal{D}	Stab(\mathcal{A}_{10}^j)
\mathcal{A}_{10}^1	5,50,52,376,374,6	I
\mathcal{A}_{10}^2	5,12,344,282,375,52	\mathbb{Z}_2
\mathcal{A}_{10}^3	25,133,40,237,378,247	\mathbb{Z}_3
\mathcal{A}_{10}^4	25,37,40,133,237,146	\mathbb{Z}_4
\mathcal{A}_{10}^5	5,12,282,327,344,304	$\mathbb{Z}_2 \times \mathbb{Z}_2$
\mathcal{A}_{10}^6	5,12,157,303,344,304	D₆



In Table 6, $\text{Stab}(\mathcal{A}_{10}^2) = \langle M_{10}^2 \rangle$, $\text{Stab}(\mathcal{A}_{10}^3) = \langle M_{10}^3 \rangle$, and $\text{Stab}(\mathcal{A}_{10}^4) = \langle M_{10}^4 \rangle$, where

$$M_{10}^2 = \begin{bmatrix} 9 & -4 & -4 \\ 9 & -3 & -6 \\ 9 & -6 & -3 \end{bmatrix}, M_{10}^3 = \begin{bmatrix} 0 & 7 & -7 \\ 1 & 7 & -7 \\ 0 & 8 & -7 \end{bmatrix} \text{ and } M_{10}^4 = \begin{bmatrix} 0 & 0 & 1 \\ 7 & 0 & 1 \\ 0 & 8 & -7 \end{bmatrix}.$$

The stabilizer group of the (10,3)-arc, \mathcal{A}_{10}^5 consists of the 4 projective matrices, each one of them has of order 2 except the identity matrix:

$$M_{10,1}^5 = I, M_{10,2}^5 = \begin{bmatrix} 0 & 2 & -1 \\ 0 & -8 & 0 \\ -7 & 3 & 0 \end{bmatrix}, M_{10,3}^5 = \begin{bmatrix} 9 & -4 & -4 \\ 9 & -3 & -6 \\ 9 & -6 & -3 \end{bmatrix} \text{ and } M_{10,4}^5 = \begin{bmatrix} -4 & 0 & 4 \\ -6 & 2 & 4 \\ -3 & 0 & 4 \end{bmatrix}.$$

The stabilizer group of the (10,3)-arc, \mathcal{A}_{10}^6 is non-abelian consisting of the 12 projective matrices, and $\text{Stab}(\mathcal{A}_{10}^6) = \langle R, S: R^6 = S^2 = I, SRS^{-1} = R^{-1} \rangle$, where

$$R = \begin{bmatrix} 0 & 2 & -1 \\ -7 & 3 & 0 \\ 0 & -8 & 0 \end{bmatrix} \text{ and } S = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$$

3.3 The classification of (v, 3)-arcs; v = 11, 12, 13, 14

In this subsection, the classification of (v, 3)-arcs, where v = 11,12,13,14, is established by classifying all such arcs up to stabilizer groups. By adding 7 preferred points

of type 3 to \mathcal{R} , we get the following (11,3)-arcs as shown in Table 7.

Table 7. Types of (11, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{11}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{11}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{11}^j)$
\mathcal{A}_{11}^1	5,6,7,8,9,10,34	I
\mathcal{A}_{11}^2	5,6,7,15,33,116,23	\mathbb{Z}_2
\mathcal{A}_{11}^3	25,133,40,237,247,6,80	\mathbb{Z}_3



\mathcal{A}_{11}^4	25,133,40,237,247,12,72	\mathbb{Z}_4
\mathcal{A}_{11}^5	5,12,344,157,303,37,228	$\mathbb{Z}_2 \times \mathbb{Z}_2$
\mathcal{A}_{11}^6	25,133,40,237,37,146,346	\mathbb{Z}_6
\mathcal{A}_{11}^7	5,6,236,224,281,31,60	\mathfrak{S}_3

The stabilizer group, $\text{Stab}(\mathcal{A}_{11}^1)$, is the trivial group. The group $\text{Stab}(\mathcal{A}_{11}^2)$ consists of two projective matrices, and hence it is isomorphic to \mathbb{Z}_2 . Further, $\text{Stab}(\mathcal{A}_{11}^3) = \langle M_{11}^3 \rangle$, $\text{Stab}(\mathcal{A}_{11}^4) = \langle M_{11}^4 \rangle$ and $\text{Stab}(\mathcal{A}_{11}^6) = \langle M_{11}^6 \rangle$, where

$$M_{11}^3 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 8 & -8 \\ 0 & 1 & -8 \end{bmatrix}, M_{11}^4 = \begin{bmatrix} 1 & 8 & -8 \\ 0 & 8 & -8 \\ 0 & 1 & -8 \end{bmatrix} \text{ and } M_{11}^6 = \begin{bmatrix} -7 & 8 & 0 \\ -7 & 8 & -1 \\ -8 & 8 & 0 \end{bmatrix}.$$

The stabilizer group of \mathcal{A}_{11}^5 has 4 projective matrices, each one of them has of order 2 except the identity matrix, say

$$M_{11,1}^5 = I, M_{11,2}^5 = \begin{bmatrix} 0 & 2 & -1 \\ 0 & -8 & 0 \\ -7 & 3 & 0 \end{bmatrix}, M_{11,3}^5 = \begin{bmatrix} 9 & -4 & -4 \\ 9 & -3 & -6 \\ 9 & -6 & -3 \end{bmatrix} \text{ and } M_{11,4}^5 = \begin{bmatrix} -4 & 0 & 4 \\ -6 & 2 & 4 \\ -3 & 0 & 4 \end{bmatrix}.$$

The group $\text{Stab}(\mathcal{A}_{11}^7)$ is non-abelian group of order 6. In fact, the generators of $\text{Stab}(\mathcal{A}_{11}^7)$ are

$$R = \begin{bmatrix} -5 & 5 & 1 \\ 6 & 5 & 0 \\ 2 & 5 & 0 \end{bmatrix} \text{ and } S = \begin{bmatrix} 0 & 0 & 1 \\ 0 & -8 & 0 \\ 7 & 0 & 0 \end{bmatrix}.$$

Adding 8 preferred points of type 3 to the arc \mathcal{R} , implies four (12,3)-arcs up to stabilizer group as shown in Table 8.

Table 8. Types of (12, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{12}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{12}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{12}^j)$
----------------------	-------------------------	-----------------------------------



\mathcal{A}_{12}^1	5,6,7,8,9,10,34	I
\mathcal{A}_{12}^2	5,6,7,15,33,116,23	\mathbb{Z}_2
\mathcal{A}_{12}^3	25,133,40,237,247,6,80	\mathbb{Z}_3
\mathcal{A}_{12}^4	5,12,344,157,303,37,228	$\mathbb{Z}_3 \times \mathbb{Z}_3$

The stabilizer groups $\text{Stab}(\mathcal{A}_{12}^2)$ and $\text{Stab}(\mathcal{A}_{12}^3)$ have 2 and 3 projective matrices, respectively. It follows that $\text{Stab}(\mathcal{A}_{12}^2) \cong \mathbb{Z}_2$ and $\text{Stab}(\mathcal{A}_{12}^3) \cong \mathbb{Z}_3$. Furthermore, The group $\text{Stab}(\mathcal{A}_{12}^4)$ is an abelian group with 9 projective matrices, and it has no element of order 9. Therefore, it is isomorphic to $\mathbb{Z}_3 \times \mathbb{Z}_3$.

Now, adding 9 preferred points of type 3 to the arc \mathcal{R} , implies four (13,3)-arcs as shown in

Table 9.

Table 9. Types of (13, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{13}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{13}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{13}^j)$
\mathcal{A}_{13}^1	5,7,15,33,45,54,59,245,315	I
\mathcal{A}_{13}^2	5,12,344,157,303,37,304,46,25	\mathbb{Z}_2
\mathcal{A}_{13}^3	5,6,7,8,72,132,133,40,146	\mathbb{Z}_3
\mathcal{A}_{13}^4	5,12,344,157,303,37,304,46,85	\mathfrak{S}_3

The stabilizer groups $\text{Stab}(\mathcal{A}_{13}^2)$ and $\text{Stab}(\mathcal{A}_{13}^3)$ have 2 and 3 projective matrices, respectively. It follows that $\text{Stab}(\mathcal{A}_{13}^2) \cong \mathbb{Z}_2$ and $\text{Stab}(\mathcal{A}_{13}^3) \cong \mathbb{Z}_3$. Furthermore, The group $\text{Stab}(\mathcal{A}_{13}^4)$ is non-abelian group with 6 projective matrices. Therefore, it is isomorphic to \mathfrak{S}_3 .



There are 3 types of (14,3)-arcs up to stabilizer group as shown in Table 10. These arcs can be constructed by adding 10 preferred points of type 3 to the arc \mathcal{R} .

Table 10. Types of (14, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{14}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{14}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{14}^j)$
\mathcal{A}_{14}^1	5,6,7,50,52,204,209,310,312,376	I
\mathcal{A}_{14}^2	5,12,157,289,298,303,304,344,356,367	\mathbb{Z}_2
\mathcal{A}_{14}^3	25,133,40,237,247,6,80,334,47,171	\mathbb{Z}_3

The stabilizer group $\text{Stab}(\mathcal{A}_{14}^1)$ is the trivial group. Further, the stabilizer groups $\text{Stab}(\mathcal{A}_{14}^2)$ and $\text{Stab}(\mathcal{A}_{14}^3)$ have 2 and 3 projective matrices, respectively. It follows that $\text{Stab}(\mathcal{A}_{14}^2) \cong \mathbb{Z}_2$ and $\text{Stab}(\mathcal{A}_{14}^3) \cong \mathbb{Z}_3$.

3.4 The classification of $(\nu, 3)$ -arcs; $\nu = 15, 16, 17, 18, 19, 20$

In this subsection, the classification of $(\nu, 3)$ -arcs, where $\nu = 15, 16, 17, 18, 19, 20$, is established by classifying all such arcs up to stabilizer groups. By adding 11 preferred points of type 3 to \mathcal{R} , we get the following (15,3)-arcs as shown in Table 11.

Table 11. Types of (15, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{15}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{15}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{15}^j)$
\mathcal{A}_{15}^1	5,6,7,15,33,45,54,59,245,315,378	I
\mathcal{A}_{15}^2	5,12,37,46,85,157,256,303,304,344,371	\mathbb{Z}_2
\mathcal{A}_{15}^3	6,25,40,47,80,133,171,237,247,315,334	$\mathbb{Z}_3 \times \mathbb{Z}_3$



All the groups in Table 11 are abelian of order 1, 2 and 9. Therefore, the stabilizer group, $\text{Stab}(\mathcal{A}_{15}^1)$, is the trivial group. The group $\text{Stab}(\mathcal{A}_{15}^2) \cong \mathbb{Z}_2$ and $\text{Stab}(\mathcal{A}_{15}^3) \cong \mathbb{Z}_3 \times \mathbb{Z}_3$.

There are 3 types of (16,3)-arcs up to stabilizer group as shown in Table 12. These arcs can be constructed by adding 12 preferred points of type 3 to the arc \mathcal{R} .

Table 12. Types of (16, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{16}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{16}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{16}^j)$
\mathcal{A}_{16}^1	5,7,15,33,45,54,59,245,315,378,316,299	I
\mathcal{A}_{16}^2	5,12,22,157,289,298,301,303,304,344,356,367	\mathbb{Z}_2
\mathcal{A}_{16}^3	6,25,40,47,80,115, 133,171,237,247,315,334	\mathbb{Z}_3

All the groups in Table 12 are abelian of order 1, 2 and 3. Therefore, the stabilizer group, $\text{Stab}(\mathcal{A}_{16}^1)$, is the trivial group. The group $\text{Stab}(\mathcal{A}_{16}^2) \cong \mathbb{Z}_2$ and $\text{Stab}(\mathcal{A}_{16}^3) \cong \mathbb{Z}_3$.

There are 3 types of (17,3)-arcs up to stabilizer group. These arcs can be established by adding 13 preferred points of type 3 to the arc \mathcal{R} . Table 13 illustrates all these arcs.

Table 13. Types of (17, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{17}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{17}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{17}^j)$
\mathcal{A}_{17}^1	5,7,15,33,45,54,59, 245,315,378,316,299,320	I
\mathcal{A}_{17}^2	5,7,15,33,45,54,59, 245,315,378,316,299,287	\mathbb{Z}_2
\mathcal{A}_{17}^3	6,25,40,47,80,115,133, 171,203,237,247,315,334	\mathbb{Z}_3

Similarly, all the groups in Table 13 are abelian of orders 1, 2 and 3. It follows that the stabilizer group, $\text{Stab}(\mathcal{A}_{17}^1)$, is the trivial group. The group $\text{Stab}(\mathcal{A}_{17}^2) \cong \mathbb{Z}_2$ and $\text{Stab}(\mathcal{A}_{17}^3) \cong \mathbb{Z}_3$.



There are 4 types of (18,3)-arcs up to stabilizer group. These arcs can be established by adding 14 preferred points of type 3 to the arc \mathcal{R} . Table 14 illustrates all these arcs.

Table 14. Types of (18, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{18}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{18}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{18}^j)$
\mathcal{A}_{18}^1	5,7,15,33,45,54,59, 245,315,378,316,299,320,254	I
\mathcal{A}_{18}^2	5,7,15,33,45,54,59, 245,315,378,316,299,287,260	\mathbb{Z}_2
\mathcal{A}_{18}^3	6,25,40,47,80,133,171, 237,247,251,315,334,355,356	\mathbb{Z}_3
\mathcal{A}_{18}^4	6,25,40,47,80,115,133, 171,203,237,247,261,315,334	$\mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_3$

The groups in Table 14 are abelian of orders 1, 2, 4 and 27. The stabilizer group, $\text{Stab}(\mathcal{A}_{18}^1)$, is the trivial group. The group $\text{Stab}(\mathcal{A}_{18}^2) \cong \mathbb{Z}_2$ and $\text{Stab}(\mathcal{A}_{18}^3) \cong \mathbb{Z}_3$. Every element in $\text{Stab}(\mathcal{A}_{18}^4)$ has order 3, so it is isomorphic to $\mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_3$.

There are 3 types of (19,3)-arcs up to stabilizer group as shown in Table 15. These arcs can be constructed by adding 15 preferred points of type 3 to the arc \mathcal{R} .

Table 15. Types of (19, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{19}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{19}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{19}^j)$
\mathcal{A}_{19}^1	5,7,15,33,45,54,59,245, 315,378,316,299,320,254,196	I
\mathcal{A}_{19}^2	5,7,15,33,45,54,59,94, 245,260,287,299,315,316,378	\mathbb{Z}_2
\mathcal{A}_{19}^3	6,25,40,47,80,115,133,171, 237,247,251,315,334,355,356	\mathbb{Z}_3

Again, all the groups in Table 15 are abelian of orders 1, 2, and 3. Hence, the stabilizer group, $\text{Stab}(\mathcal{A}_{19}^1)$, is the trivial group, and the group $\text{Stab}(\mathcal{A}_{19}^2) \cong \mathbb{Z}_2$ and $\text{Stab}(\mathcal{A}_{19}^3) \cong \mathbb{Z}_3$.



There are 2 types of (20,3)-arcs up to stabilizer group as shown in Table 16. These arcs can be constructed by adding 16 preferred points of type 3 to the arc \mathcal{R} .

Table 16. Types of (20, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{20}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{20}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{20}^j)$
\mathcal{A}_{20}^1	5,7,15,33,45,54,59,245, 315,378,316,299,320,254,196,244	I
\mathcal{A}_{20}^2	6,25,40,47,80,115,133,171, 203,237,247,251,315,334,355,356	\mathbb{Z}_3

All the groups in Table 16 are abelian of orders 1 and 3. Hence, the stabilizer group, $\text{Stab}(\mathcal{A}_{20}^1)$, is the trivial group, and the group, $\text{Stab}(\mathcal{A}_{20}^2) \cong \mathbb{Z}_3$.

3.5 The classification of $(\nu, 3)$ -arcs; $\nu = 21, 22, 23, 24, 25, 26, 27, 28, 29, 30$

In this subsection, we establish the distinct $(\nu, 3)$ -arcs up to their stabilizer groups, where $\nu = 21, \dots, 30$. By similar arguments in the previous subsections, if we adding 17 preferred points of type 3 to the arc \mathcal{R} , we get the following (21,3)-arcs as shown in Table 17.

Table 17. Types of (21, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{21}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{21}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{21}^j)$
\mathcal{A}_{21}^1	5,7,15,33,45,54,59,245,315, 378,316,299,320,254,196,244,309	I
\mathcal{A}_{21}^2	6,25,40,47,80,115,133,171,203, 237,247,251,261,315,334,355,356	\mathbb{Z}_3
\mathcal{A}_{21}^3	12,16,38,43,66,110,171,217,253, 282,291,298,327,344,349,367,378	D₁₈



\mathcal{A}_{21}^4	12,16,38,43,66,110,171,217,253, 282,291,298,327,344,349,367,381	\mathbf{D}_{20}
----------------------	--	-------------------

The stabilizer group, $\text{Stab}(\mathcal{A}_{21}^1)$, is the trivial group, and the group $\text{Stab}(\mathcal{A}_{21}^2) \cong \mathbb{Z}_3$ because it has three projective matrices. The stabilizer group of the (21,3)-arc, \mathcal{A}_{21}^3 is non-abelian consisting of the 36 projective matrices, and $\text{Stab}(\mathcal{A}_{21}^3) = \langle R, S: R^{18} = S^2 = I, SRS^{-1} = R^{-1} \rangle$, where

$$R = \begin{bmatrix} -4 & 9 & -4 \\ -3 & -3 & 6 \\ -6 & -2 & 8 \end{bmatrix} \text{ and } S = \begin{bmatrix} 0 & 6 & -5 \\ 9 & 0 & -3 \\ 0 & 0 & 4 \end{bmatrix}.$$

Furthermore, The stabilizer group of the (21,3)-arc, \mathcal{A}_{21}^4 is non-abelian consisting of the 40 projective matrices, and $\text{Stab}(\mathcal{A}_{21}^4) = \langle R, S: R^{20} = S^2 = I, SRS^{-1} = R^{-1} \rangle$, where

$$R = \begin{bmatrix} 8 & 7 & 5 \\ 9 & -1 & -9 \\ -8 & 5 & -3 \end{bmatrix} \text{ and } S = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 9 & 0 \\ 5 & 0 & 0 \end{bmatrix}.$$

There are 4 types of (22,3)-arcs up to stabilizer group as shown in Table 18. These arcs can be constructed by adding 18 preferred points of type 3 to the arc \mathcal{R} .

Table 18. Types of (22, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{22}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{22}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{22}^j)$
\mathcal{A}_{22}^1	12,16,38,43,66,110,111,217,253,282,291,298,327,344,349,367,5,368	\mathbf{I}
\mathcal{A}_{22}^2	12,16,38,43,66,110,111,217,253,282,291,298,327,344,349,367,5,171	\mathbb{Z}_2
\mathcal{A}_{22}^3	12,16,38,43,66,110,171,217,253,282,291,298,327,344,349,367,378,373	$\mathbb{Z}_2 \times \mathbb{Z}_2$
\mathcal{A}_{22}^4	12,16,38,39,43,66,110,171,217,253,282,291,298,327,344,349,367,378	\mathbf{D}_4

In Table 18, $\text{Stab}(\mathcal{A}_{22}^2)$ is isomorphic to \mathbb{Z}_2 because it has two projective matrices. The group $\text{Stab}(\mathcal{A}_{22}^3)$ has 4 projective matrices, each one of them has order 2 except the identity, so it is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$. Furthermore, The stabilizer group of the (22,3)-arc, \mathcal{A}_{22}^4 is non-abelian

consisting of the 8 projective matrices, and $\text{Stab}(\mathcal{A}_{22}^4) = \langle R, S: R^4 = S^2 = I, SRS^{-1} = R^{-1} \rangle$, where

$$R = \begin{bmatrix} -3 & 5 & -1 \\ 2 & 6 & -8 \\ -9 & 7 & 2 \end{bmatrix} \text{ and } S = \begin{bmatrix} 0 & -1 & 2 \\ 0 & 8 & 0 \\ -6 & 4 & 0 \end{bmatrix}.$$

There are 4 types of (23,3)-arcs up to stabilizer group as shown in Table 19. These arcs can be constructed by adding 19 preferred points of type 3 to the arc \mathcal{R} .

Table 19. Types of (23, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{23}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{23}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{23}^j)$
\mathcal{A}_{23}^1	12,16,38,43,66,110,111,217,253,282,291, 298,327,344,349,367,5,368,171	I
\mathcal{A}_{23}^2	12,16,38,43,66,110,171,217,253,282,291, 298,327,344,349,367,378,373,361	\mathbb{Z}_2
\mathcal{A}_{23}^3	12,16,38,39,43,66,75,110,171,217,253,282,291, 298,327,344,349,367,378	$\mathbb{Z}_2 \times \mathbb{Z}_2$
\mathcal{A}_{23}^4	5,12,16,38,43,66,110,111,156,171,217,253,282,291, 298,327,344,349,367	\mathfrak{S}_3

In Table 19, $\text{Stab}(\mathcal{A}_{23}^2)$ is isomorphic to \mathbb{Z}_2 because it has two projective matrices. The group $\text{Stab}(\mathcal{A}_{23}^3)$ has 4 projective matrices, each one of them has order 2 except the identity, so it is



isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$. Furthermore, The stabilizer group of the (23,3)-arc, \mathcal{A}_{23}^4 is non-abelian with 6 projective matrices, and $\text{Stab}(\mathcal{A}_{23}^4) = \langle R, S \rangle$, where

$$R = \begin{bmatrix} 6 & -3 & -2 \\ 9 & -6 & 0 \\ -5 & -2 & 0 \end{bmatrix} \text{ and } S = \begin{bmatrix} 9 & 0 & -8 \\ 5 & -5 & 8 \\ 7 & 0 & -9 \end{bmatrix}.$$

By similar arguments, if we adding 20 preferred points of type 3 to the arc \mathcal{R} , we get the following (24,3)-arcs as shown in Table 20.

Table 20. Types of (24, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{24}^j = \mathcal{R} \cup \mathcal{D}$

\mathcal{A}_{24}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{24}^j)$
\mathcal{A}_{24}^1	12,16,38,43,66,110,171, 217,253,282,291, 298,327,344, 349,367,378,373,361,226	I
\mathcal{A}_{24}^2	12,16,28,38,39,43,66, 75,110,171,217,253,282,291, 298,327,344,349,367,378	\mathbb{Z}_2
\mathcal{A}_{24}^3	12,16,38,43,66,102,110, 171,217,253,282,291, 298,327, 344,349,361,367,373,378	$\mathbb{Z}_2 \times \mathbb{Z}_2$

In Table 20, $\text{Stab}(\mathcal{A}_{24}^2)$ is isomorphic to \mathbb{Z}_2 because it has two projective matrices. The group $\text{Stab}(\mathcal{A}_{24}^3)$ has 4 projective matrices, each one of them has order 2 except the identity, so it is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$.

Again, if we add 21 preferred points of type 3 to the arc \mathcal{R} , we get the three types of (25,3)-arcs up to stabilizer group as shown in Table 21.

Table 21. Types of (25, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ; $\mathcal{A}_{25}^j = \mathcal{R} \cup \mathcal{D}$



\mathcal{A}_{25}^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_{25}^j)$
\mathcal{A}_{25}^1	12,16,38,43,66,110,171, 217,253,282,291, 298,327,344, 349,367,378,373,361,226	I
\mathcal{A}_{25}^2	12,16,28,38,39,43,66, 75,110,171,217,253,282,291, 298,327,344,349,367,378	\mathbb{Z}_2
\mathcal{A}_{25}^3	12,16,38,43,66,102,110, 171,217,253,282,291, 298,327, 344,349,361,367,373,378	$\mathbb{Z}_2 \times \mathbb{Z}_2$

In Table 21, $\text{Stab}(\mathcal{A}_{25}^2)$ is isomorphic to \mathbb{Z}_2 because it has two projective matrices. The group $\text{Stab}(\mathcal{A}_{25}^3)$ has 4 projective matrices, each one of them has order 2 except the identity, so it is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$.

Now, adding 22, 23, 24 and 25 preferred points of type 3 to the arc \mathcal{R} , will give us two (26,3)-arcs, three (27,3)-arcs, two (28,3)-arcs and two (29,3)-arcs, respectively as shown in Table 22.

Table 22. Types of (26, 3), (27, 3), (28, 3), (29, 3)-arcs up to stabilizer groups in \mathbb{P}_{19}^2 ;

$$\mathcal{A}_k^j = \mathcal{R} \cup \mathcal{D}, k = 26, 27, 28, 29$$

\mathcal{A}_k^j	Points of \mathcal{D}	$\text{Stab}(\mathcal{A}_k^j)$
\mathcal{A}_{26}^1	12,16,38,43,66,110,111,217,253,282,291, 298,327,344,349,367,5,368,319,171,39,337	I
\mathcal{A}_{26}^2	12,16,28,38,39,43,66,75,108,110,171,217,253,282,291,	\mathbb{Z}_2



	296,298,327,344,349,367,378	
\mathcal{A}_{27}^1	12,16,38,43,66,110,111,217,253,282,291, 298,327,344,349,367,5,379,314,266,286,274,171	I
\mathcal{A}_{27}^2	12,16,28,38,39,43,66,75,108,110,171,217,253,282,291, 296,298,322,327,344,349,367,378	\mathbb{Z}_2
\mathcal{A}_{27}^3	12,16,28,38,39,43,66,75,108,110,171,217,253,282,291, 296,298,310,327,344,349,367,378	$\mathbb{Z}_2 \times \mathbb{Z}_2$
\mathcal{A}_{28}^1	12,16,38,43,66,110,111,217,253,282,291, 298,327,344,349,367,5,379,314,266,286,274,279,171	I
\mathcal{A}_{28}^2	12,16,38,43,66,110,111,217,253,282,291, 298,327,344,349,367,5,379,314,266,286,274,351,171	\mathbb{Z}_2
\mathcal{A}_{29}^1	12,16,38,43,66,110,111,217,253,282,291, 298,327,344,349,367,5,379,314,266,171,155,274,279,286	I
\mathcal{A}_{29}^2	12,16,38,43,66,110,111,217,253,282,291, 298,327,344,349,367,5,379,314,266,286,274,279,171,351	\mathbb{Z}_2

In Table 22, The stabilizer groups, $\text{Stab}(\mathcal{A}_{26}^1)$, $\text{Stab}(\mathcal{A}_{27}^1)$, $\text{Stab}(\mathcal{A}_{28}^1)$ and $\text{Stab}(\mathcal{A}_{29}^1)$ are the trivial groups, because they have only one projective matrix. The stabilizer groups, $\text{Stab}(\mathcal{A}_{26}^2)$, $\text{Stab}(\mathcal{A}_{27}^2)$, $\text{Stab}(\mathcal{A}_{28}^2)$ and $\text{Stab}(\mathcal{A}_{29}^2)$ are isomorphic to \mathbb{Z}_2 because it has two projective matrices. The group $\text{Stab}(\mathcal{A}_{27}^3)$ has 4 projective matrices, each one of them has order 2 except the identity, so it is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$.

Finally, if we add 26 preferred points of type 3 to the arc \mathcal{R} , we have

only (30,3)-arc, say $\mathcal{A}_{30} = \mathcal{R} \cup \mathcal{D}$, where



$$\mathcal{D} = \left\{ \begin{array}{l} 5,9,12,16,17,38,43,66,110,171,189,217,253,254, \\ 282,284,291,298,307,318,327,344,349,367,381,358 \end{array} \right\}$$

The stabilizer group of the $(30,3)$ -arc, \mathcal{A}_{30} is isomorphic to \mathbb{Z}_2 . In fact, $\text{Stab}(\mathcal{A}_{30}) = \langle M \rangle$, where

$$M = \begin{bmatrix} 6 & 6 & 8 \\ -7 & -9 & 9 \\ 4 & 7 & -8 \end{bmatrix}.$$

According to the arguments in sections 3.1, 3.2, 3.3, 3.4 and 3.5, we present the following theorem:

Theorem. A In the projective plane of order 19, \mathbb{P}_{19}^2 , we have:

1. There are 9 projectively inequivalent $(7,3)$ -arcs up to stabilizer groups.
2. There are 8 projectively inequivalent $(6,3)$ -arcs up to stabilizer groups.
3. For $k = 7,11$, there are 7 projectively inequivalent $(k, 3)$ -arcs up to stabilizer groups.
4. For $k = 8,10$, there are 6 projectively inequivalent $(k, 3)$ -arcs up to stabilizer groups.
5. For $k = 5,12,13,18,21,22,23$, there are 4 projectively inequivalent $(k, 3)$ -arcs up to stabilizer groups.
6. For $k = 14,15,16,17,19,24,25,27$, there are 3 projectively inequivalent $(k, 3)$ -arcs up to stabilizer groups.
7. For $k = 20,26,28,29$, there are 2 projectively inequivalent $(k, 3)$ -arcs up to stabilizer groups.
8. There is only one projectively inequivalent $(30,3)$ -arcs up to stabilizer groups.

Conclusions

In summary, the large size of complete $(k, 3)$ -arcs which is 30 is constructed by classify all

$(k, 3)$ -arcs according to their stabilizer groups. Additionally, there are two projectively inequivalent $(k, 3)$ -arcs for an integer k , where $k = 20,26,28,29$. Moreover, we find three projectively inequivalent $(k, 3)$ -arcs up to stabilizer groups for the value of k , where $k = 14,15,16,17,19,24,25,27$. As well as, for $k = 8,10$, there are six projectively inequivalent $(k, 3)$ -arcs up to stabilizer groups. Furthermore, there are seven types of projectively inequivalent $(k, 3)$ -arcs up to stabilizer groups for $k = 7,11$.

xxope (FESEM).



References

- [1] J. W. P. Hirschfeld, J. A. Thas, J. A., General Galois geometries, Oxford: Clarendon Press, 1378(1991), <https://doi.org/10.1007/978-1-4471-6790-7>
- [2] G. R. Cook, Arcs in a finite projective plane (Doctoral dissertation, University of Sussex), 2011.
- [3] S. Marcugini, A. Milani, F. Pambianco, Maximal $(n, 3)$ -arcs in $PG(2, 11)$, Discrete Math., 208(1999) 421-426, [https://doi.org/10.1016/s0012-365x\(99\)00202-2](https://doi.org/10.1016/s0012-365x(99)00202-2)
- [4] S. Marcugini, A. Milani, F. Pambianco, Maximal $(n, 3)$ -arcs in $PG(2, 13)$, Discrete Math., 294(1-2)(2005) 139-145, <https://doi.org/10.1007/s00022-004-1777-4>
- [5] J.W.P. Hirschfeld, E.V.D. Pichanick, Bounds for arcs of arbitrary degree in finite Desarguesian planes, J. Comb. Des., 24(4)(2016) 184-196, <https://doi.org/10.1002/jcd.21426>
- [6] K. Coolsaet, H. Sticker, A full classification of the complete k -arcs of $PG(2, 23)$ and $PG(2, 25)$, J. Comb. Des., 17(6)(2009) 459-477, <https://doi.org/10.1002/jcd.20211>
- [7] K. Coolsaet, H. Sticker, The complete $(k, 3)$ -arcs of $PG(2, q)$, $q \leq 13$. J. Comb. Des., 20(2)(2012) 89-111, <https://doi.org/10.1002/jcd.20293>
- [8] H.J. Al-Mayyahi, M.A. Alabbod, Lower Bound for $m_3(2, 37)$ and Related Code, Turk. J. Comput. Math. Educ, 12(14)(2021)959-969, <https://doi.org/10.17762/turcomat.v12i14.10377>
- [9] H.J. Al-Mayyahi, M.A. Alabbod, On the $(4n, 3)$ -arcs in $PG(2, q)$ and the related linear codes, Int. J. Nonlinear Anal. Appl, 12(2)(2021)2589-2599. <http://dx.doi.org/10.22075/ijnaa.2021.5430>.
- [10] R.H. Hanoon, M.A. Ibrahim, Large (k, n) -arcs in the Projective Plane of Order 37, Bas. J. Sci., 41.2(2023) 178-197, <https://doi.org/10.29072/basjs.20230201>
- [11] M. Ibrahim, Islam, Large $(k, 3)$ -arcs in $PG(2, 19)$ and the related linear codes. J. Kufa math. comput., 11(1)(2024) 43–54, <https://doi.org/10.31642/JoKMC/2018/110108>
- [12] S. Ball, and A. Blokhuis, The classification of maximal arcs in small Desarguesian planes, Bull. Belg. Math. Soc. Simon Stevin., 9(3)(2002).433-445, <https://doi.org/10.36045/bbms/1102715068>
- [13] D.J.S. Robinson, Fundamental Concepts of Group Theory. In: A Course in the Theory of Groups. Graduate Texts in Mathematics, Springer, New York, NY, vol 80 (1996), https://doi.org/10.1007/978-1-4419-8594-1_1



- [14] J. Cameron, Introduction to Algebra, Oxford, online edn, Oxford Academic, 1997, <https://doi.org/10.1093/oso/9780198569138.001.0001>.
- [15] J. W. P. Hirschfeld, Projective geometries over finite fields. Oxford University Press. 1998, <https://doi.org/10.1093/oso/9780198502951.001.0001>
- [16] D. Bartoli, A., Davydov A., G. Faina, S. Marcugini, and F. Pambianco, 2012. On sizes of complete arcs in PG (2, q). Discrete Math., 312(3)(2012)680-698. <https://doi.org/10.1016/j.disc.2011.07.002>
- [17] S. Linton, GAP, groups, algorithms, programming. ACM Commun. in Comput. Algebra, 41(3)(2007)108-109, <https://doi.org/10.1145/1358190.1358201>

تصنيف أقواس الدرجة الثالثة في $PG(2,19)$ حسب زمر التثبيت

اسلام عباس عبد الزهرة ، محمد علي ابراهيم

قسم الرياضيات - كلية العلوم - جامعة البصرة- البصرة - العراق

البريد الإلكتروني: mohammed.ibrahim@uobasrah.edu.iq

المستخلص

القوس من النمط $(n, 3)$ ، \mathcal{K} في المستوى الإسقاطي $PG(2, q)$ ذو الحجم n والدرجة الثالثة عبارة عن مجموعة من النقاط n بحيث يلتقي بها كل خط في المستوى في أقل من أو يساوي ثلاث نقاط، وكذلك القوس \mathcal{K} تكون كاملة إذا لم تكن موجودة في قوس من النمط $(n + 1, 3)$. في هذا البحث تم تقديم تصنيف الأقواس من الدرجة الثالثة في $PG(2, q)$ بالتفصيل حسب زمر التثبيت الخاصة بها. الدافع للعمل في المستوى الإسقاطي للطلب 19 ذو شقين. أولاً، حجم القوس الأكبر $(n, 3)$ غير معروف. ثانياً، عدد الأقواس $(n, 3)$ أعلى بكثير في المستوى الإسقاطي من الرتبة 19 مما هو عليه في المستوى الإسقاطي من الرتبة q لاجل $q < 19$ ، مما يعطي عدداً كبيراً من الأقواس ذو النمط $(n, 3)$ للدراسة.

